



## King's Research Portal

DOI:

[10.1111/ene.13847](https://doi.org/10.1111/ene.13847)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Ray Chaudhuri, K., Antonini, A., Robieson, W. Z., Sanchez-Solino, O., Bergmann, L., Poewe, W., & on behalf of the GLORIA study co-investigators (2019). Burden of nonmotor symptoms in Parkinson's disease patients predicts improvement in quality of life during treatment with levodopacarbidoopa intestinal gel. *European Journal of Neurology*, 26(4), 581-e43. <https://doi.org/10.1111/ene.13847>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# IN OUR WORLD, A STEADY FLOW KEEPS PATIENTS ALIVE.

LET'S TAKE CARE OF IT.  
TOGETHER.

## Driving Excellence in Neurology.

Be part of one of humankind's most ambitious projects to help uncover the workings of the nervous system. EAN connects 45,000 specialists in neurology in 47 countries across Europe. Promoting excellence, cutting-edge science and innovative therapies for a better life for more than 430 million patients in Europe alone. Read more on [ean.org](http://ean.org)

Great Minds.





# Burden of non-motor symptoms in Parkinson's disease patients predicts improvement in quality of life during treatment with levodopa-carbidopa intestinal gel

K. Ray Chaudhuri<sup>a,b</sup> , A. Antonini<sup>c</sup> , W. Z. Robieson<sup>d</sup>, O. Sanchez-Solano<sup>d</sup>, L. Bergmann<sup>d</sup> and W. Poewe<sup>e</sup> on behalf of the GLORIA study co-investigators

<sup>a</sup>Parkinson Foundation International Centre of Excellence, King's College Hospital, Denmark Hill, London; <sup>b</sup>Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK. <sup>c</sup>Department of Neuroscience, University of Padua, Padua, Italy.

<sup>d</sup>AbbVie Inc., North Chicago, IL, USA. and <sup>e</sup>Medical University of Innsbruck, Innsbruck, Austria

## Keywords:

levodopa-carbidopa intestinal gel, non-motor symptoms, Parkinson's disease, quality of life, routine patient care

Received 7 April 2018  
Accepted 12 October 2018

*European Journal of Neurology* 2019, **26**: 581–587,  
e42–e43

doi:10.1111/ene.13847

**Background and purpose:** GLORIA, a registry conducted with 375 advanced Parkinson's disease patients treated with levodopa-carbidopa intestinal gel (LCIG) for 24 months in routine clinical care, demonstrated significant reductions from baseline in 'off' time and 'on' time with dyskinesia and improvements in the Non-Motor Symptom Scale (NMSS) total and individual domain scores, and in Parkinson's Disease Questionnaire 8 item (PDQ-8) total score.

**Methods:** Associations between baseline NMSS burden (NMSB), the multi-domain NMSS total score and the PDQ-8 total score were investigated for 233 patients. Baseline NMSB was assigned to five numerical categories defined by the NMSS total cutoff scores (0–20, 21–40, 41–60, 61–80 and >80). Pearson and Spearman correlations were calculated at month 24.

**Results:** The response of LCIG was assessed using validated criteria after 24 months. The proportion of patients decreasing  $\geq 30$  NMSS score points was 47% in the most affected NMSB category (NMSS total score > 80). A positive association was noted between baseline NMSB and NMSS total score (0.57,  $P < 0.0001$ ), as well as between NMSS total score and PDQ-8 total score (0.46,  $P < 0.0001$ ). Associations between improvements of the NMSS domain sleep/fatigue and PDQ-8 total score (0.32,  $P = 0.0001$ ) as well as between the NMSS domain mood/cognition and PDQ-8 total score (0.37,  $P < 0.0001$ ) were also shown.

**Conclusions:** This analysis demonstrated positive associations between NMSS baseline burden and improvements of non-motor symptoms. Improvements of non-motor symptoms were associated with improved quality of life in advanced parkinsonian patients during a 2-year treatment with LCIG and reflect the long-term non-motor efficacy of this treatment.

## Introduction

Parkinson's disease (PD) is characterized by a combination of motor and non-motor symptoms (NMSs) underpinned by relevant neuropathological changes

[1–7] which contribute significantly to morbidity, loss of autonomy, institutionalization and increases in healthcare costs [8,9]. The Non-Motor Symptom Scale (NMSS) and the NMS questionnaire are validated tools [10,11] to assess a broad range of NMSs and have been used in many recent PD studies [12–18].

Management of PD requires recognition of both motor and non-motor disturbances as well as an understanding of the relationship between these symptoms and how they can be ameliorated by medications [9,16]. Treatment of NMSs can be challenging as these symptoms are often underestimated or

Correspondence: K. Ray Chaudhuri, Department of Basic and Clinical Neuroscience, The Maurice Wohl Clinical Neuroscience Institute, King's College London, Cutcombe Road, London SE5 9RT, UK (tel.: +44 203 299 7154; fax: +44 208 772 0669; e-mail: Ray.chaudhuri@kcl.ac.uk).

[The copyright line for this article was changed on 18 January 2019 after original online publication]

unrecognized by clinicians and remain untreated [2]. When identified, there is a common perception that dopaminergic drugs are not efficacious in improving these symptoms [9,19]. Furthermore, the progression of NMSs contributes importantly to the decline in quality of life (QoL) in patients with PD [9,11,16,20]. There are emerging clinical data that the overall burden of NMSs seems to be the major determinant of QoL, and that NMSs, as a whole, have a greater impact than motor symptoms on QoL in patients with motor fluctuations [21]. A good body of evidence suggests that motor symptoms and NMSs originate from distinct pathophysiological pathways although NMSs may respond to dopaminergic treatment [22,23].

A longitudinal multivariate analysis study showed that the total NMS burden (NMSB) significantly predicted the QoL scores whilst motor scores did not. The burden of NMSs, in particular sleep, mood and attention, had a significant impact on the QoL of PD patients [24].

The comprehensive clinical data of the GLORIA long-term registry in 375 patients with advanced PD [25] offered the opportunity to investigate whether baseline (BL) NMSB would predict improvements of NMSs as well as improvements between NMSs and QoL following a 24-month levodopa-carbidopa intestinal gel (LCIG) treatment.

## Patients and methods

### Study design

GLORIA, a multinational, non-interventional, observational registry, was conducted at 75 specialized movement disorder centres across 18 countries in advanced PD patients with persistent motor complications. The study protocol was approved by national and/or local independent ethics committees and health authorities at each participating institution and country. All patients provided written informed consent before enrolment in the registry. The results of the GLORIA registry, both the 12-month interim and the final 24-month analysis, were published [15,26].

### Patients

Male and female patients with advanced PD and severe motor complications eligible for LCIG treatment based on the approved European Commission Summary of Product Characteristics and national reimbursement criteria/local pathways were enrolled in the GLORIA study. Clinical observations of the initial 24-month LCIG treatment phase were collected.

### Efficacy and safety

The following efficacy and safety outcomes were assessed: Unified Parkinson's Disease Rating Scale (UPDRS) parts II, III, IV and V; complications of therapy [UPDRS IV items 32 and 39 modified according to the Movement Disorder Society–UPDRS (corresponding parts 4.3 and 4.1) to allow for calculation of actual hours of 'off' time and 'on' time with dyskinesias]. NMSs were assessed using the NMSS. Patient-reported QoL was assessed using the short-form Parkinson's Disease Questionnaire (PDQ-8) [27] and generic EuroQoL-5 Dimensions quality of life instrument questionnaire (EQ-5D). All adverse drug reactions (ADR) were recorded during LCIG treatment. ADRs were defined as those adverse events that in the opinion of the investigator had at least a reasonable possibility of being causally related to the study drug, as described previously [15,26]. Clinical data were recorded at BL prior to initiation of LCIG using a temporal nasojejunal tube, at discharge from the hospital following percutaneous endoscopic gastrostomy with jejunal extension tube placement (day 1), and at months 6, 12, 18 and 24.

### Post hoc analyses

Potential associations between NMSB at BL and improvements of NMSs as well as between improvements of NMSs and QoL during LCIG treatment were assessed. The following five categories of BL NMSB were numerically defined: NMSS total score 0–20, 21–40, 41–60, 61–80 and >80. Correlation analysis was performed between categories of BL NMSB and improvements of NMSS total score, improvements of NMSS total score and PDQ-8 total score, and improvements of NMSS domain scores (sleep/fatigue and mood/cognition) and PDQ-8 total score. The analyses are presented by Pearson and Spearman correlation coefficients (prob > |r| under H<sub>0</sub>:  $\rho = 0$ ) and shown in regression plots.

## Results

All patients with a BL NMSS assessment (233 out of 375 patients included in GLORIA) were included in this *post hoc* analysis. 12- and 24-month NMSS recordings were available for 194 and 170, respectively, out of 258 patients who completed GLORIA. The mean  $\pm$  SD NMSS total score, demographics and PD characteristics at BL of patients allocated to the five numerically defined NMSB categories (0–20, 21–40, 41–60, 61–80 and >80) and the total *post hoc* analysis population (PHP) are shown in Table 1.

12.0% of the patients were allocated to the lowest NMSB category (NMSS total score 0–20,  $10.8 \pm 4.77$ ) and 38.2% to the highest NMSB category (NMSS total score >80;  $112.8 \pm 27.38$ ). The mean age ( $66.2 \pm 8.52$  years in the PHP) and time since PD diagnosis ( $12.5 \pm 5.93$  years in the PHP) were similar, and time with dyskinesia ( $4.2 \pm 3.66$  h in the PHP) and ‘off’ time ( $5.9 \pm 3.02$  h in the PHP) were comparable across all five NMSB categories, whilst the PDQ-8 total scores ( $46.8 \pm 18.63$  in the PHP) increased from  $35.9 \pm 17.47$  in the lowest to  $54.3 \pm 18.81$  in the highest NMSB category.

The median NMSS and PDQ-8 total scores for the five NMSB categories and the complete PHP at BL, 12 months and 24 months are presented in Fig. 1. The median NMSS total score changes showed a trend toward more substantial improvements from the lowest to the higher BL burden categories, at similar magnitudes at 12 months and 24 months, being highest in the >80 NMSB category. The median of the PDQ-8 total score changes did not show a consistent trend across all five categories, and improvements showed a comparable magnitude at 12 months and 24 months. (Fig. 1).

The NMSS responder rates by BL NMSB categories are shown in Fig. 2. The rates for the overall response (improvement  $\geq 5$  score points) increased gradually from the lowest (0–20) to the higher NMSB categories and reached a maximum of 58% in the highest category (>80). Also, the responder rates for improvements of  $\geq 10$ ,  $\geq 20$  and  $\geq 30$  score points increased gradually from the lowest (0–20) to the highest NMSS total score category (>80) and reached a maximum response rate of 47% for improvements of  $\geq 30$  in NMSS scores.

The results of the correlation analysis are shown in Table 2. The Pearson correlation coefficient between

the BL NMSS total score and the NMSS total score improvements ( $0.57$ ,  $P < 0.0001$ ) is strong. The Pearson correlation coefficient between the improvements of the NMSS total score and the PDQ-8 total score improvements was moderate ( $0.46$ ,  $P < 0.0001$ ). The Pearson correlation coefficient between improvements of the NMSS domain sleep/fatigue and the NMSS domain mood/cognition and the PDQ-8 total score improvements was moderate ( $0.32$ ,  $P = 0.0001$ ;  $0.37$ ,  $P < 0.0001$ , respectively). The corresponding regression plots are presented in Fig. 3.

The ADRs have been described in the interim and final publication of the GLORIA registry [25,26] and did not show any new findings compared to the known safety profile of LCIG.

## Discussion


The data from this study provide us with novel insights into the possible effect of levodopa on NMSB assessed by a validated tool in PD. This *post hoc* analysis of the GLORIA study revealed for the first time that LCIG improved NMSs across all NMSB categories, even in those with very severe NMSs at BL, as well as QoL. Most importantly, it showed that the best predictor of improvement was BL NMS severity. These novel findings complement previous literature on the predication of QoL by NMSB [11,24]. In addition, the data are supportive of positron emission tomography imaging data suggesting a correlation of synaptic dopamine release and improvement of NMSs in PD [25].

The analysis of the GLORIA database included assessments of motor and non-motor symptoms and QoL with results comparable to a number of recent studies [11,15–18,20,22,24]. The results of this first,


**Table 1** Demographics and baseline disease characteristics in patients allocated to one of the five baseline NMSB categories defined by the NMSS total score cut-off scores 0–20 ( $N = 28$ ), 21–40 ( $N = 41$ ), 41–60 ( $N = 39$ ), 61–80 ( $N = 36$ ) and >80 ( $N = 89$ ) and in the total *post hoc* analysis population ( $N = 233$ )

	Baseline NMSS total score burden					
	0–20 ( $N = 28$ )	21–40 ( $N = 41$ )	41–60 ( $N = 39$ )	61–80 ( $N = 36$ )	>80 ( $N = 89$ )	All ( $N = 233$ )
Age (years)	$64.6 \pm 8.98$	$64.9 \pm 9.12$	$66.4 \pm 7.97$	$67.5 \pm 9.10$	$66.6 \pm 8.13$	$66.2 \pm 8.52$
BMI	$24.8 \pm 4.17$	$25.1 \pm 4.40$	$26.5 \pm 5.35$	$24.7 \pm 4.10$	$25.2 \pm 4.23$	$25.2 \pm 4.40$
Time since PD diagnosis (years)	$13.4 \pm 5.45$	$12.7 \pm 4.47$	$11.7 \pm 5.96$	$12.3 \pm 7.58$	$12.6 \pm 5.97$	$12.5 \pm 5.93$
UPDRS part IV modified item 32: Time with dyskinesia (h)	$3.0 \pm 2.45$	$3.8 \pm 2.94$	$4.8 \pm 3.50$	$5.3 \pm 4.71$	$4.2 \pm 3.86$	$4.2 \pm 3.66$
UPDRS part IV modified item 39: ‘Off’ time (h)	$6.0 \pm 2.32$	$5.7 \pm 3.47$	$5.2 \pm 2.69$	$4.9 \pm 3.27$	$6.6 \pm 2.87$	$5.9 \pm 3.02$
<b>NMSS total score</b>	<b><math>10.8 \pm 4.77</math></b>	<b><math>29.6 \pm 6.41</math></b>	<b><math>51.9 \pm 6.32</math></b>	<b><math>70.5 \pm 5.17</math></b>	<b><math>112.8 \pm 27.38</math></b>	<b><math>69.18 \pm 42.13</math></b>
Total PDQ-8 score	$35.9 \pm 17.47$	$38.6 \pm 16.79$	$45.2 \pm 18.98$	$46.3 \pm 12.33$	$54.3 \pm 18.81$	$46.8 \pm 18.63$

BMI, body mass index; NMSB, NMSS burden; NMSS, Non-Motor Symptom Scale; PDQ-8, Parkinson’s Disease Questionnaire 8 item; UPDRS, Unified Parkinson’s Disease Rating Scale. Values are presented as mean  $\pm$  SD or percentage of patients. NMSS total scores in bold represents the key NMS data reflecting rising NMS burden as pre cut off scores in top panel.



**Figure 1** (a) Median NMSS total score and (b) median PDQ-8 total at BL, and median change to BL scores (absolute change and percent of change) in the five BL NMSB categories defined by the BL NMSS total score ranges 0–20 ( $N = 28$ ), 21–40,  $N = 41$ ), 41–60 ( $N = 39$ ), 61–80 ( $N = 36$ ) and >80 ( $N = 89$ ) at month 12 and month 24 in the total *post hoc* analysis population ( $N = 233$ ). [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**Figure 2** NMSS responder rates (NMSS total score improvements  $\geq 5$  score points,  $\geq 10$  score points,  $\geq 20$  score points and  $\geq 30$  score points) by the BL NMSB categories and the total *post hoc* analysis population (PHP). [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

**Table 2** Correlation coefficients between (a) the BL NMSS total score burden and the NMSS total score improvement, (b) NMSS total score improvements and PDQ-8 total score improvements, (c) NMSS subdomain sleep/fatigue improvement and PDQ-8 total score improvement and (d) NMSS subdomain mood/cognition improvement and PDQ-8 total score improvement

Variable	N	Mean	SD	Median	Minimum	Maximum
(a) Baseline NMSS total score burden and NMSS total score improvement						
NMS total score improvement	170	16.7	43.17	18	−122	148
NMS total score at BL	233	69.2	42.13	66	2	217
<b>Pearson correlation coefficient</b>						
Correlation coefficient		0.56515				0.56879
P value		<0.0001				<0.0001
N		170	N			170
(b) NMSS total score improvement and PDQ-8 total score improvement						
NMS total score improvement	170	16.7	43.17	18	−122	148
PDQ-8 total score improvement	152	7.1	21.00	6.25	−53.125	65.625
<b>Pearson correlation coefficient</b>						
Correlation coefficient		0.46392				0.43782
P value		<0.0001				<0.0001
N		140	N			140
(c) NMSS domain sleep/fatigue improvement and PDQ-8 total score improvement						
NMS domain sleep/fatigue improvement	167	5.3	11.08	3	−20	36
PDQ-8 total score improvement	152	7.1	21.00	6.25	−53.125	65.625
<b>Pearson correlation coefficient</b>						
Correlation coefficient		0.32101				0.31003
P value		0.0001				0.0002
N		139	N			139
(d) NMSS domain mood/cognition improvement and PDQ-8 total score improvement						
NMS domain mood/cognition improvement	167	3.1	12.62	1	−30	48
PDQ-8 total score improvement	152	7.1	21.00	6.25	−53.125	65.625
<b>Pearson correlation coefficient</b>						
Correlation coefficient		0.36793				0.38388
P value		<0.0001				<0.0001
N		139	N			139

Method: prob > |r| under H0:  $\rho = 0$ .

large multinational, long-term registry demonstrated sustained improvements with LCIG of motor and non-motor symptoms and in NMSS subdomains (particularly sleep/fatigue, mood/cognition and gastrointestinal domains), as well as QoL in advanced PD patients at 12 and 24 months [25,26].

These *post hoc* analyses provide evidence for (i) the rationale of dopaminergic treatment for NMSs of PD in conjunction with continued motor improvements as has been postulated previously [19,28]; (ii) an association between BL NMSB and treatment-related improvements of NMSs; and (iii) a moderate correlation between improvement of NMSS total score and improvement of QoL as measured by PDQ-8 in advanced PD patients.


Similar improvements of NMSs and QoL were demonstrated in two other open-label studies with smaller patient collectives over a 12-month and a mean 48-month follow-up period [29,30].

Many NMSs occur early in PD and in *de novo* PD (some may even predate the diagnosis of PD currently) [6,31] whilst NMSs dominate through the natural history of PD [23]. The NMSB rises as the

condition progresses. Given ours was an advanced PD cohort, it is not surprising that only 12% in this cohort with advanced PD showed mild to moderate NMSB (0–20). On the other hand, the largest proportion of patients (38%) were in the most severe NMSB category (>80). An association of higher NMSS total scores with worse QoL was shown [16], and the NMSB was reported to be the best predictor of QoL [11]. Whilst the NMSs decreased in all subgroups carrying mild to severe NMSB, the reduction of NMSs was most pronounced in the most severe NMSB category (−43 total NMSS score points). The PDQ-8 total score decreased in parallel with all NMSB subgroups at month 24. Similar results were published in two other studies, one cross-sectional [11] and one longitudinal, both showing that NMSB significantly predicted QoL scores [24].

A possible threshold for a minimal clinically important difference (MCID) of the NMSS has been discussed by many authors based on a multitude of randomized, comparative and open-label clinical trials which have used the NMSS as an outcome measure [32–34]. Currently, since a minimally important





**Figure 3** Regression plots (a) between the BL NMSS total score burden and the NMSS total score improvements; (b) between the NMSS total score improvements and the PDQ-8 total score improvements; (c) between the improvements of the NMSS sleep/fatigue domain and the PDQ-8 total score; (d) between the improvements of the NMSS mood/cognition domain and the PDQ-8 total score. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

difference is not fully established and not agreed upon by experts, a spectrum of numerical responder cutoffs was chosen, capturing both mild and strong responders. Even with the strictest responder definition (decrease of  $\geq 30$  NMSS score points), the proportion of responders was 47% in the most affected NMSB category (NMSS total score  $> 80$ ). Martinez-Martin *et al.* proposed an NMSS MCID of 13.91 [35]. Thus, patients with a decrease of  $\geq 20$  NMSS score points in our study could be considered responders. Similarly, Horváth *et al.* suggested a PDQ-8 MCID of  $-5.94$  [36], and reductions of median PDQ-8 scores were above this threshold in the NMSB categories 41–60, 61–80 and  $> 80$ .

The predictive nature of BL NMSB was confirmed by a strong correlation between the NMSS total BL score and the NMSS total score improvements ( $0.57$ ,  $P < 0.0001$ ). Improvement of overall NMSs may be partially due to reduction of dopamine-related non-motor fluctuations [20].

There is now compelling evidence (clinical and statistical) that the total NMSB may be the key determinant of QoL, especially in advanced PD [28]. In our study, the Pearson correlation coefficient between NMSS total score improvement and improvement of the PDQ-8 total score at month 24 showed a moderate correlation ( $0.46$ ,  $P < 0.0001$ ).

The NMSS allows for the nine subdomains to be tested individually as has been performed in previous studies [11,18], and specifically it was of interest to explore the effect of LCIG on sleep/fatigue as well as mood/cognition domains based on evidence obtained from EuroInf [18] and other studies. In our study, a significant association existed between improvements of the two NMSS subdomain scores (sleep/fatigue and mood/cognition) and the PDQ-8 total score improvements ( $< 0.0001$ ), confirming this observation from the other open-label studies. Similar observations were reported in another study: the NMSS domains sleep/fatigue, mood/apathy and attention/memory were most significantly predictive of QoL change [24]. Sleep in PD is a complex phenomenon driven by various neurotransmitter system abnormalities [26]. However, a considerable element of sleep dysfunction in PD is dopaminergic and consequently in theory responsive to levodopa therapy. Our data support this observation



and also form a basis for delivery of personalised medicine for PD, a key unmet need, with LCIG [37].

In conclusion, GLORIA results demonstrate that LCIG decreased NMSB, with largest benefits observed in those patients with most severe BL NMSB, and confirmed the significant association with QoL improvements.

### Acknowledgements

Medical writing support was provided by Urs E. Gasser, ClinResearch, funded by AbbVie, and the statistical analysis was performed by Christoph Meyenberg, Koehler eClinical, funded by AbbVie.

### Disclosure of conflicts of interest

Dr Chaudhuri was a study investigator and has received honorarium from UCB, AbbVie, Britannia, Mundipharma, Boehringer Ingelheim and GSK Pharmaceuticals for lecturing at symposia. He has acted as a consultant for UCB, AbbVie, Britannia, Neuronova and Mundipharma. He has received research funding from Parkinson's UK, NIHR, PDNMG, as well as educational grants from UCB, Britannia, AbbVie, GSK Pharmaceuticals, Boehringer Ingelheim and Neuronova. Dr Chaudhuri receives royalties from Oxford University Press and holds intellectual

property rights for the Kings Parkinson's Pain Scale and Parkinson's Disease Sleep Scale 2. Dr Chaudhuri acknowledges independent research part funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London, and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author and not necessarily those of the NHS, the NIHR or the Department of Health, UK. A. Antonini was a study investigator and has received compensation for consultancy and speaker-related activities from UCG, Boston Scientific, Boehringer Ingelheim, AbbVie and Zambon. He has received research support from Mundipharma. W. Poewe was a study investigator and has received compensation from AbbVie, AstraZeneca, Teva, Novartis, GSK, Boehringer Ingelheim, UCB, Orion Pharma, Zambon and Merz Pharmaceuticals (consultancy and lecture fees in relation to clinical drug development programmes for PD) outside the submitted work. He has received royalties from Thieme, Wiley Blackwell and Oxford University Press. W.Z. Robieson, Olga Sanchez-Solino and L. Bergmann are employees of AbbVie and may hold AbbVie stock and/or stock options. AbbVie contributed to the design, study conduct and financial support for the study. AbbVie participated in the interpretation of data, review and approval of the publication.

## References

- Jankovic J. Parkinson's disease: clinical features and diagnosis. *J Neurol Neurosurg Psychiatry* 2008; **79**: 368–376.
- Antonini A, Chaudhuri KR, Martinez-Martin P, Odin P. Oral and infusion levodopa-based strategies for managing motor complications in patients with Parkinson's disease. *CNS Drugs* 2010; **24**: 119–129.
- Kalia LV, Lang AE. Parkinson disease. *Lancet Neurol* 2015; **386**: 896–912.
- Chapuis S, Ouchchane L, Metz O, Gerbaud L, Durif F. Impact of the motor complications of Parkinson's disease on the quality of life. *Mov Disord* 2005; **20**: 224–230.
- Chaudhuri KR, Odin P, Antonini A, Martinez-Martin P. Parkinson's disease: the non-motor issues. *Parkinsonism Relat Disord* 2011; **17**: 717–723.
- Titova N, Padmakumar C, Lewis SJG, Chaudhuri KR. Parkinson's: a syndrome rather than a disease? *J Neural Transm* 2017; **124**: 907–914.
- Jellinger K. Neuropathobiology of non-motor symptoms in Parkinson disease. *J Neural Transm (Vienna)* 2015; **122**: 1429–1440.
- Lundqvist C, Beiske AG, Reiertsen O, Kristiansen IS. Real life cost and quality of life associated with continuous intraduodenal levodopa infusion compared with oral treatment in Parkinson patients. *J Neurol* 2014; **261**: 2438–2445.
- Palhagen SE, Sydow O, Johansson A, et al. Levodopa-carbidopa intestinal gel (LCIG) treatment in routine care of patients with advanced Parkinson's disease: an open-label prospective observational study of effectiveness, tolerability and healthcare costs. *Parkinsonism Relat Disord* 2016; **29**: 17–23.
- Chaudhuri KR, Martinez-Martin P, Brown RG. The metric properties of a novel non-motor symptoms scale for Parkinson's disease: results from an international pilot study. *Mov Disord* 2007; **22**: 1901–1911.
- Martinez-Martin P, Rodriguez-Blazquez C, Kurtis MM, Chaudhuri KR, NMSS Validation Group. The impact of non-motor symptoms on health-related quality of life of patients with Parkinson's disease on behalf of the NMSS Validation Group. *Mov Disord* 2011; **26**: 399–406.
- Eggert K, Schrader C, Hahn M. Continuous jejunal levodopa infusion in patients with advanced Parkinson disease: practical aspects and outcome of motor and non-motor complications. *Clin Neuropharmacol* 2008; **31**: 151–166.
- Honig H, Antonini A, Martinez-Martin P. Intrajejunal levodopa infusion in Parkinson's disease: a pilot multicenter study of effects on non-motor symptoms and quality of life. *Mov Disord* 2009; **24**: 1468–1474.
- Reddy P, Martinez-Martin A, Rizos P, et al. Intrajejunal levodopa versus conventional therapy in Parkinson disease: motor and nonmotor effects. *Clin Neuropharmacol* 2012; **35**: 205–207.
- Antonini A, Yegin A, Preda C, Bergmann L, Poewe W, on behalf of the GLORIA study investigators and coordinators. Global long-term study on motor and non-motor symptoms and safety of levodopa-carbidopa intestinal gel in routine care of advanced Parkinson's disease patients; 12-month interim outcomes. *Parkinsonism Relat Disord* 2015; **21**: 231–235.
- Santos-García D, de la Fuente-Fernández R. Impact of non-motor symptoms on health-related and perceived quality of life in Parkinson's disease. *J Neurol Sci* 2013; **332**: 136–140.
- Antonini A, Bauer L, Dohin E, et al. Effects of rotigotine transdermal patch in patients with Parkinson's disease presenting with non-motor symptoms – results of a double-blind, randomized, placebo-controlled trial. *Eur J Neurol* 2015; **22**: 1400–1407.
- Martinez-Martin P, Reddy P, Katzenschlager R, et al. EuroInf: a multicenter comparative observational study of apomorphine and levodopa infusion in Parkinson's disease. *Mov Disord* 2015; **30**: 510–516.
- Chaudhuri KR, Schapira AH. Non-motor symptoms of Parkinson's disease: dopaminergic pathophysiology and treatment. *Lancet Neurol* 2009; **8**: 464–474.
- Antonini A, Barone P, Marconi R, et al. The progression of non-motor symptoms in Parkinson's disease and their contribution to motor disability and quality of life. *J Neurol* 2012; **259**: 2621–2631.
- Buck P, Trautman H, Clark J. Scales for assessing non-motor symptom severity changes in Parkinson's disease patients with symptom fluctuations. *Int J Neurosci* 2010; **120**: 523–530.
- Swick TJ, Friedman JH, Chaudhuri KR, et al. Associations between severity of motor function and nonmotor symptoms in Parkinson's disease: a *post hoc* analysis of the RECOVER study. *Eur Neurol* 2014; **71**: 140–147.
- Schapira A, Chaudhuri KR, Jenner P. Non-motor features of Parkinson disease. *Nat Rev Neurosci* 2017; **18**: 435–450.
- Prakash KM, Nadkarni NV, Lye WK, Yong MH, Tan EK. The impact of non-motor symptoms on the quality of life of Parkinson's disease patients: a longitudinal study. *Eur J Neurol* 2016; **23**: 854–860.
- Politis M, Sauerbier A, Loane C, et al. Sustained striatal dopamine levels following intestinal levodopa/infusions in Parkinson's disease patients. *Mov Disord* 2017; **32**: 235–240.
- Antonini A, Poewe W, Chaudhuri KR, et al., on behalf of the GLORIA study co-investigators. Levodopa-carbidopa intestinal gel in advanced Parkinson's: final results of the GLORIA registry. *Parkinsonism Relat Disord* 2017; **45**: 13–20.
- Jenkinson C, Fitzpatrick R, Peto V, Greenhall R, Hyman N. The PDQ-8: development and validation of a short-form Parkinson's disease questionnaire. *Psychol Health* 1997; **12**: 805–814.
- Chaudhuri KR, Odin P, Antonini A, Martinez-Martin P. Parkinson's disease: the non-motor issues. *Parkinsonism Relat Disord* 2011; **17**: 717–723.
- Juhász A, Aschermann Z, Ács P, et al. Levodopa/carbidopa intestinal gel can improve both motor and non-motor experiences of daily living in Parkinson's disease: an open-label study. *Parkinsonism Relat Disord* 2017; **37**: 79–86.
- Bohlega S, Abou Al-Shaar H, Alkhairallah T, Al-Ajlan F, Hasan N, Alkahtani K. Levodopa-carbidopa intestinal gel infusion therapy in advanced Parkinson's disease: single Middle Eastern center experience. *Eur Neurol* 2015; **74**: 227–236.

31. Chaudhuri KR, Naidu Y. Early Parkinson's disease and non-motor issues. *J Neurol* 2008; **255**: 33–38.
32. Martin P, Rodriguez-Blazquez C, Abe K, *et al.* International study on the psychometric attributes of the non-motor symptoms scale in Parkinson disease. *Neurology* 2009; **73**: 1584–1591.
33. Sloan JA, Cella D, Hays RD. Clinical significance of patient-reported questionnaire data: another step toward consensus. *J Clin Epidemiol* 2005; **58**: 1217–1219.
34. Crosby RD, Kolotkin RL, Williams GR. Defining clinically meaningful change in health-related quality of life. *J Clin Epidemiol* 2003; **56**: 395–407.
35. Martinez-Martin P, Rodriguez-Blazquez C, Abe K, *et al.* International study on the psychometric attributes of the Non-Motor Symptoms Scale in Parkinson disease. *Neurology* 2009; **73**: 1584–1591.
36. Horváth K, Aschermann Z, Kovács M, *et al.* Changes in quality of life in Parkinson's disease: how large must they be to be relevant? *Neuroepidemiology* 2017; **48**: 1–8. (ISSN: 1423-0208).
37. Titova N, Chaudhuri KR. Personalized medicine and nonmotor symptoms in Parkinson's Disease. *Int Rev Neurobiol* 2017; **134**: 1257–1281.